

## Study Guide

**Factoring Trinomials**

To factor a trinomial of the form  $ax^2 + bx + c$ , follow Example 1 below.

**Example 1:** Factor  $2d^2 + 15d + 18$ .  
The product of 2 and 18 is 36.  
You need to find two integers whose *product* is 36 and whose *sum* is 15.

Factors of 36	Sum of Factors
1, 36	$1 + 36 = 37$
2, 18	$2 + 18 = 20$
3, 12	$3 + 12 = 15$

$$\begin{aligned}
 2d^2 + 15d + 18 &= 2d^2 + (12 + 3)d + 18 \\
 &= 2d^2 + 12d + 3d + 18 \\
 &= (2d^2 + 12d) + (3d + 18) \\
 &= 2d(d + 6) + 3(d + 6) && \text{Factor the GCF from each group.} \\
 &= (2d + 3)(d + 6) && \text{Use the distributive property.}
 \end{aligned}$$

To factor a trinomial of the form given above when  $a = 1$ , you need to find only the factors of  $c$  whose sum is  $b$ .

**Example 2:** Factor  $x^2 + 7x + 10$ .

Since 2 and 5 are factors of 10 whose sum is 7,  
 $x^2 + 7x + 10 = (x + 2)(x + 5)$ .

The same pattern can be used to factor a trinomial  $ax^2 + bx + c$  when  $a = 1$  and  $c$  is negative. When this occurs, the factors of the trinomial are a *difference* and a *sum*.

**Complete.**

1.  $x^2 - 5x - 14 = (x + \underline{\quad})(x - 7)$

2.  $a^2 + 13a + 36 = (a + 9)(a \underline{\quad} 4)$

3.  $p^2 - 25 = (p + 5)(p \underline{\quad} \underline{\quad})$

4.  $x^2 - 6xy - 16y^2 = (x \underline{\quad} \underline{\quad})(x + 2y)$

5.  $49 - n^2 = (7 \underline{\quad} \underline{\quad})(\underline{\quad} \underline{\quad} n)$

6.  $a^4 + 3xa^2 - 10x^2 = (a^2 - \underline{\quad})(\underline{\quad} \underline{\quad} 5x)$

**Factor each trinomial, if possible. If the trinomial cannot be factored using integers, write prime.**

7.  $x^2 + 12y + 32$

8.  $x^2 - x - 6$

9.  $x^2 - 4x - 21$

10.  $y^2 + 22y + 121$

11.  $9 - 7n + n^2$

12.  $a^2 - 16a + 64$

13.  $3x^2 + 2x - 8$

14.  $18h^2 - 27h - 5$

15.  $28x^2 + 60x - 25$

16.  $48x^2 + 22x - 15$

17.  $-4y^2 + 19y - 21$

18.  $6a^2 - 7a + 18$